



US005202413A

United States Patent [19]

[11] Patent Number: **5,202,413**

Spinu

[45] Date of Patent: **Apr. 13, 1993**

[54] **ALTERNATING (ABA)_n POLYLACTIDE
BLOCK COPOLYMERS**

[75] Inventor: **Maria Spinu, Wilmington, Del.**

[73] Assignee: **E. I. Du Pont de Nemours and
Company, Wilmington, Del.**

[21] Appl. No.: **816,201**

[22] Filed: **Dec. 31, 1991**

[51] Int. Cl.⁵ **C08G 63/08**

[52] U.S. Cl. **528/354; 525/415;
528/26; 528/28; 528/76; 528/83; 528/300;
528/301; 528/341**

[58] Field of Search **528/28, 26, 76, 83,
528/300, 301, 341, 354; 525/415**

[56] **References Cited**

U.S. PATENT DOCUMENTS

2,878,236	3/1959	Young et al. .	
3,169,945	2/1965	Hostettler et al. .	
4,328,322	5/1982	Baroni	528/68 X
4,699,974	10/1987	Evans	528/354
4,950,735	8/1990	Vanderbilt et al.	528/354
5,028,667	7/1991	McLain et al.	525/415

FOREIGN PATENT DOCUMENTS

5060290	1/1990	Australia .
3911557	8/1989	Fed. Rep. of Germany .
59-27923	2/1984	Japan .

OTHER PUBLICATIONS

Barskaya et al., "Polymer Science USSR," (translated from) vol. 25, No. 7, pp. 1788-1793 (1983).

Feng et al., "J. Polym. Sci: Letters Ed." vol. 21 pp. 593-600 (1983).

Song et al., "Macromolecules" vol. 17 pp. 2764-2767 (1984).

Grijpma et al., "Polymer Bulletin" vol. 25 pp. 327-333 (1991).

Hamitou et al., "J. Polym. Sci: Pol. Chem. Ed." vol. 15, pp. 1035-1041 (1977).

K. J. Ivin, T. Saegusa, "Ring Opening Polymerization" vol. 1, Ch. 7 pp. 461-521 (1984).

Primary Examiner—Nathan M. Nutter

Assistant Examiner—Shelley Wright

Attorney, Agent, or Firm—Robert B. Stevenson

[57] **ABSTRACT**

Polymers of the general formula R—(A-B-A-I)_xA-B-A-R and their preparation are disclosed. In the general formula, -A- is a polyhydroxy acid residue preferably a polymer of either 98 to 100% L-lactide or 98 to 100% D-lactide having a molecular weight of 500 to 40,000 and preferably 2,000 to 20,000, -B- is an oligomeric diol or diamine residue, having a number average molecular weight of 500 to 20,000 and preferably 2,000 to 8,000. The polymers are prepared by polymerizing a lactone onto the diol or diamine followed by reacting the resulting polymeric H-A-B-A-H with an diisocyanate or diacyl chloride to form the -L- chain extending units in the general formula.

25 Claims, No Drawings